

FACT SHEET FOR HEALTHCARE PROVIDERS ADMINISTERING VACCINE (VACCINATION PROVIDERS)

EMERGENCY USE AUTHORIZATION (EUA) OF THE PFIZER-BIONTECH COVID-19 VACCINE TO PREVENT CORONAVIRUS DISEASE 2019 (COVID-19)

The U.S. Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) to permit the emergency use of the unapproved product, **Pfizer-BioNTech COVID-19 Vaccine**, for active immunization to prevent COVID-19 in individuals 12 years of age and older.

SUMMARY OF INSTRUCTIONS FOR COVID-19 VACCINATION PROVIDERS

Vaccination providers enrolled in the federal COVID-19 Vaccination Program must report all vaccine administration errors, all serious adverse events, cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and cases of COVID-19 that result in hospitalization or death following administration of Pfizer-BioNTech COVID-19 Vaccine. See “MANDATORY REQUIREMENTS FOR PFIZER-BIONTECH COVID-19 VACCINE ADMINISTRATION UNDER EMERGENCY USE AUTHORIZATION” for reporting requirements.

The Pfizer-BioNTech COVID-19 Vaccine is a suspension for intramuscular injection administered as a series of two doses (0.3 mL each) 3 weeks apart.

See this Fact Sheet for instructions for preparation and administration. This Fact Sheet may have been updated. For the most recent Fact Sheet, please see www.cvdvaccine.com.

For information on clinical trials that are testing the use of the Pfizer-BioNTech COVID-19 Vaccine for active immunization against COVID-19, please see www.clinicaltrials.gov.

DESCRIPTION OF COVID-19

Coronavirus disease 2019 (COVID-19) is an infectious disease caused by the novel coronavirus, SARS-CoV-2, that appeared in late 2019. It is predominantly a respiratory illness that can affect other organs. People with COVID-19 have reported a wide range of symptoms, ranging from mild symptoms to severe illness. Symptoms may appear 2 to 14 days after exposure to the virus. Symptoms may include: fever or chills; cough; shortness of breath; fatigue; muscle or body aches; headache; new loss of taste or smell; sore throat; congestion or runny nose; nausea or vomiting; diarrhea.

DOSAGE AND ADMINISTRATION

Storage and Handling

During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.

Do not refreeze thawed vials.

Frozen Vials Prior to Use

Cartons of Pfizer-BioNTech COVID-19 Vaccine Multiple Dose Vials arrive in thermal containers with dry ice. Once received, remove the vial cartons immediately from the thermal container and preferably store in an ultra-low temperature freezer between -80°C to -60°C (-112°F to -76°F) until the expiry date printed on the label. Alternatively, vials may be stored at -25°C to -15°C (-13°F to 5°F) for up to 2 weeks. Vials must be kept frozen and protected from light until ready to use. Vials stored at -25°C to -15°C (-13°F to 5°F) for up to 2 weeks may be returned one time to the recommended storage condition of -80°C to -60°C (-112°F to -76°F). Total cumulative time the vials are stored at -25°C to -15°C (-13°F to 5°F) should be tracked and should not exceed 2 weeks.

If an ultra-low temperature freezer is not available, the thermal container in which the Pfizer-BioNTech COVID-19 Vaccine arrives may be used as temporary storage when consistently re-filled to the top of the container with dry ice. Refer to the re-icing guidelines packed in the original thermal container for instructions regarding the use of the thermal container for temporary storage. The thermal container maintains a temperature range of -90°C to -60°C (-130°F to -76°F). Storage of the vials between -96°C to -60°C (-141°F to -76°F) is not considered an excursion from the recommended storage condition.

Transportation of Frozen Vials

If local redistribution is needed and full cartons containing vials cannot be transported at -90°C to -60°C (-130°F to -76°F), vials may be transported at -25°C to -15°C (-13°F to 5°F). Any hours used for transport at -25°C to -15°C (-13°F to 5°F) count against the 2-week limit for storage at -25°C to -15°C (-13°F to 5°F). Frozen vials transported at -25°C to -15°C (-13°F to 5°F) may be returned one time to the recommended storage condition of -80°C to -60°C (-112°F to -76°F).

Thawed Vials Before Dilution

Thawed Under Refrigeration

Thaw and then store undiluted vials in the refrigerator [2°C to 8°C (35°F to 46°F)] for up to 5 days (120 hours). A carton of 25 vials or 195 vials may take up to 2 or 3 hours, respectively, to thaw in the refrigerator, whereas a fewer number of vials will thaw in less time.

Thawed at Room Temperature

For immediate use, thaw undiluted vials at room temperature [up to 25°C (77°F)] for 30 minutes. Thawed vials can be handled in room light conditions. Vials must reach room temperature before dilution.

Undiluted vials may be stored at room temperature for no more than 2 hours.

Transportation of Thawed Vials

Available data support transportation of one or more thawed vials at 2°C to 8°C (35°F to 46°F) for up to 12 hours. Any hours used for transport at 2°C to 8°C (35°F to 46°F) count against the 120-hour limit for storage at 2°C to 8°C (35°F to 46°F).

Vials After Dilution

- After dilution, store vials between 2°C to 25°C (35°F to 77°F) and use within 6 hours from the time of dilution.
- During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.
- Any vaccine remaining in vials must be discarded after 6 hours.
- Do not refreeze.

Dosing and Schedule

The Pfizer-BioNTech COVID-19 Vaccine is administered intramuscularly as a series of two doses (0.3 mL each) 3 weeks apart.

There are no data available on the interchangeability of the Pfizer-BioNTech COVID-19 Vaccine with other COVID-19 vaccines to complete the vaccination series. Individuals who have received one dose of Pfizer-BioNTech COVID-19 Vaccine should receive a second dose of Pfizer-BioNTech COVID-19 Vaccine to complete the vaccination series.

Dose Preparation

Prior to Dilution

- The Pfizer-BioNTech COVID-19 Vaccine Multiple Dose Vial contains a volume of 0.45 mL, supplied as a frozen suspension that does not contain preservative. Each vial must be thawed and diluted prior to administration.
- Vials may be thawed in the refrigerator [2°C to 8°C (35°F to 46°F)] or at room temperature [up to 25°C (77°F)] (*see Storage and Handling*).
- Refer to thawing instructions in the panels below.



Dilution

Dilute the vial contents using 1.8 mL of 0.9% Sodium Chloride Injection, USP (not provided) to form the Pfizer-BioNTech COVID-19 Vaccine. ONLY use 0.9% Sodium Chloride Injection, USP as the diluent. This diluent is not packaged with the vaccine


and must be sourced separately. Do not use bacteriostatic 0.9% Sodium Chloride Injection or any other diluent. Do not add more than 1.8 mL of diluent.

After dilution, one vial contains 6 doses of 0.3 mL. Vial labels and cartons may state that after dilution, a vial contains 5 doses of 0.3 mL. The information in this Fact Sheet regarding the number of doses per vial after dilution supersedes the number of doses stated on vial labels and cartons.


- Refer to dilution and dose preparation instructions in the panels below.

THAWING PRIOR TO DILUTION	
 <p>No more than 2 hours at room temperature (up to 25 °C / 77 °F)</p>	<ul style="list-style-type: none"> • Thaw vial(s) of Pfizer-BioNTech COVID-19 Vaccine before use either by: <ul style="list-style-type: none"> ○ Allowing vial(s) to thaw in the refrigerator [2°C to 8°C (35°F to 46°F)]. A carton of vials may take up to 3 hours to thaw, and thawed vials can be stored in the refrigerator for up to five days (120 hours). ○ Allowing vial(s) to sit at room temperature [up to 25°C (77°F)] for 30 minutes. • Using either thawing method, vials must reach room temperature before dilution and must be diluted within 2 hours.
 <p>Gently x 10</p>	<ul style="list-style-type: none"> • Before dilution invert vaccine vial gently 10 times. • <u>Do not shake.</u> • Inspect the liquid in the vial prior to dilution. The liquid is a white to off-white suspension and may contain white to off-white opaque amorphous particles. • Do not use if liquid is discolored or if other particles are observed.


DILUTION




- Obtain sterile 0.9% Sodium Chloride Injection, USP. Use only this as the diluent.
- Using aseptic technique, withdraw 1.8 mL of diluent into a transfer syringe (21-gauge or narrower needle).
- Cleanse the vaccine vial stopper with a single-use antiseptic swab.
- Add 1.8 mL of 0.9% Sodium Chloride Injection, USP into the vaccine vial.




- Equalize vial pressure before removing the needle from the vial by withdrawing 1.8 mL air into the empty diluent syringe.



- Gently invert the vial containing the Pfizer-BioNTech COVID-19 Vaccine 10 times to mix.
- Do not shake.
- Inspect the vaccine in the vial.
- The vaccine will be an off-white suspension. Do not use if vaccine is discolored or contains particulate matter.

	<ul style="list-style-type: none"> • Record the date and time of dilution on the Pfizer-BioNTech COVID-19 Vaccine vial label. • Store between 2°C to 25°C (35°F to 77°F). • Discard any unused vaccine 6 hours after dilution.
---	---

PREPARATION OF INDIVIDUAL 0.3 mL DOSES OF PFIZER-BIONTECH COVID-19 VACCINE

	<ul style="list-style-type: none"> • Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab, and withdraw <u>0.3 mL</u> of the Pfizer-BioNTech COVID-19 Vaccine preferentially using a low dead-volume syringe and/or needle. • Each dose must contain 0.3 mL of vaccine. • If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and any excess volume. • Administer immediately.
--	---

Administration

Visually inspect each dose in the dosing syringe prior to administration. The vaccine will be an off-white suspension. During the visual inspection,

- verify the final dosing volume of 0.3 mL.
- confirm there are no particulates and that no discoloration is observed.
- do not administer if vaccine is discolored or contains particulate matter.

Administer the Pfizer-BioNTech COVID-19 Vaccine intramuscularly.

After dilution, vials of Pfizer-BioNTech COVID-19 Vaccine contain six doses of 0.3 mL of vaccine. Low dead-volume syringes and/or needles can be used to extract six doses from a single vial. If standard syringes and needles are used, there may not be sufficient volume to extract a sixth dose from a single vial. Irrespective of the type of syringe and needle:

- Each dose must contain 0.3 mL of vaccine.
- If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and content.
- Do not pool excess vaccine from multiple vials.

Contraindications

Do not administer Pfizer-BioNTech COVID-19 Vaccine to individuals with known history of a severe allergic reaction (e.g., anaphylaxis) to any component of the Pfizer-BioNTech COVID-19 Vaccine (*see Full EUA Prescribing Information*).

Warnings

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of Pfizer-BioNTech COVID-19 Vaccine.

Monitor Pfizer-BioNTech COVID-19 Vaccine recipients for the occurrence of immediate adverse reactions according to the Centers for Disease Control and Prevention (CDC) guidelines (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-anaphylaxis.html>).

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the Pfizer-BioNTech COVID-19 Vaccine.

Pfizer-BioNTech COVID-19 Vaccine may not protect all vaccine recipients.

Adverse Reactions

Adverse Reactions in Clinical Trials

Adverse reactions following the Pfizer-BioNTech COVID-19 Vaccine that have been reported in clinical trials include injection site pain, fatigue, headache, muscle pain, chills, joint pain, fever, injection site swelling, injection site redness, nausea, malaise, and lymphadenopathy (*see Full EUA Prescribing Information*).

Adverse Reactions in Post Authorization Experience

Severe allergic reactions, including anaphylaxis, and other hypersensitivity reactions (e.g., rash, pruritus, urticaria, angioedema), diarrhea, vomiting, and pain in extremity (arm) have been reported following administration of the Pfizer-BioNTech COVID-19 Vaccine outside of clinical trials.

Additional adverse reactions, some of which may be serious, may become apparent with more widespread use of the Pfizer-BioNTech COVID-19 Vaccine.

Use with Other Vaccines

There is no information on the co-administration of the Pfizer-BioNTech COVID-19 Vaccine with other vaccines.

INFORMATION TO PROVIDE TO VACCINE RECIPIENTS/CAREGIVERS

As the vaccination provider, you must communicate to the recipient or their caregiver, information consistent with the “Fact Sheet for Recipients and Caregivers” (and provide a copy or direct the individual to the website www.cvdvaccine.com to obtain the Fact Sheet) prior to the individual receiving each dose of Pfizer-BioNTech COVID-19 Vaccine, including:

- FDA has authorized the emergency use of the Pfizer-BioNTech COVID-19 Vaccine, which is not an FDA-approved vaccine.
- The recipient or their caregiver has the option to accept or refuse Pfizer-BioNTech COVID-19 Vaccine.
- The significant known and potential risks and benefits of Pfizer-BioNTech COVID-19 Vaccine, and the extent to which such risks and benefits are unknown.
- Information about available alternative vaccines and the risks and benefits of those alternatives.

For information on clinical trials that are testing the use of the Pfizer-BioNTech COVID-19 Vaccine to prevent COVID-19, please see www.clinicaltrials.gov.

Provide a vaccination card to the recipient or their caregiver with the date when the recipient needs to return for the second dose of Pfizer-BioNTech COVID-19 Vaccine.

Provide the v-safe information sheet to vaccine recipients/caregivers and encourage vaccine recipients to participate in v-safe. V-safe is a new voluntary smartphone-based tool that uses text messaging and web surveys to check in with people who have been vaccinated to identify potential side effects after COVID-19 vaccination. V-safe asks questions that help CDC monitor the safety of COVID-19 vaccines. V-safe also provides second-dose reminders if needed and live telephone follow-up by CDC if participants report a significant health impact following COVID-19 vaccination. For more information, visit: www.cdc.gov/vsafe.

MANDATORY REQUIREMENTS FOR PFIZER-BIONTECH COVID-19 VACCINE ADMINISTRATION UNDER EMERGENCY USE AUTHORIZATION

In order to mitigate the risks of using this unapproved product under EUA and to optimize the potential benefit of Pfizer-BioNTech COVID-19 Vaccine, the following items are required. Use of unapproved Pfizer-BioNTech COVID-19 Vaccine for active immunization to prevent COVID-19 under this EUA is limited to the following (all requirements **must** be met):

1. Pfizer-BioNTech COVID-19 Vaccine is authorized for use in individuals 12 years of age and older.
2. The vaccination provider must communicate to the individual receiving the Pfizer-BioNTech COVID-19 Vaccine or their caregiver, information consistent with the “Fact Sheet for Recipients and Caregivers” prior to the individual receiving Pfizer-BioNTech COVID-19 Vaccine.
3. The vaccination provider must include vaccination information in the state/local jurisdiction’s Immunization Information System (IIS) or other designated system.
4. The vaccination provider is responsible for mandatory reporting of the following to the Vaccine Adverse Event Reporting System (VAERS):
 - vaccine administration errors whether or not associated with an adverse event,
 - serious adverse events* (irrespective of attribution to vaccination),
 - cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and
 - cases of COVID-19 that result in hospitalization or death.

Complete and submit reports to VAERS online at <https://vaers.hhs.gov/reportevent.html>. For further assistance with reporting to VAERS call 1-800-822-7967. The reports should include the words “Pfizer-BioNTech COVID-19 Vaccine EUA” in the description section of the report.

5. The vaccination provider is responsible for responding to FDA requests for information about vaccine administration errors, adverse events, cases of MIS in adults and children, and cases of COVID-19 that result in hospitalization or death following administration of Pfizer-BioNTech COVID-19 Vaccine to recipients.

* Serious adverse events are defined as:

- Death;
- A life-threatening adverse event;
- Inpatient hospitalization or prolongation of existing hospitalization;
- A persistent or significant incapacity or substantial disruption of the ability to conduct normal life functions;
- A congenital anomaly/birth defect;
- An important medical event that based on appropriate medical judgement may jeopardize the individual and may require medical or surgical intervention to prevent one of the outcomes listed above.

OTHER ADVERSE EVENT REPORTING TO VAERS AND PFIZER INC.

Vaccination providers may report to VAERS other adverse events that are not required to be reported using the contact information above.


To the extent feasible, report adverse events to Pfizer Inc. using the contact information below or by providing a copy of the VAERS form to Pfizer Inc.

Website	Fax number	Telephone number
www.pfizersafetyreporting.com	1-866-635-8337	1-800-438-1985

ADDITIONAL INFORMATION

For general questions, visit the website or call the telephone number provided below.

To access the most recent Pfizer-BioNTech COVID-19 Vaccine Fact Sheets, please scan the QR code provided below.

Global website	Telephone number
www.cvdvaccine.com 	1-877-829-2619 (1-877-VAX-CO19)

AVAILABLE ALTERNATIVES

There is no approved alternative vaccine to prevent COVID-19. There may be clinical trials or availability under EUA of other COVID-19 vaccines.

FEDERAL COVID-19 VACCINATION PROGRAM

This vaccine is being made available for emergency use exclusively through the CDC COVID-19 Vaccination Program (the Vaccination Program). Healthcare providers must enroll as providers in the Vaccination Program and comply with the provider requirements. Vaccination providers may not charge any fee for the vaccine and may not charge the vaccine recipient any out-of-pocket charge for administration. However, vaccination providers may seek appropriate reimbursement from a program or plan that covers COVID-19 vaccine administration fees for the vaccine recipient (private insurance, Medicare, Medicaid, Health Resources & Services Administration [HRSA] COVID-19 Uninsured Program for non-insured recipients). For information regarding provider requirements and enrollment in the CDC COVID-19 Vaccination Program, see <https://www.cdc.gov/vaccines/covid-19/provider-enrollment.html>.

Individuals becoming aware of any potential violations of the CDC COVID-19 Vaccination Program requirements are encouraged to report them to the Office of the Inspector General, U.S. Department of Health and Human Services, at 1-800-HHS-TIPS or <https://TIPS.HHS.GOV>.

AUTHORITY FOR ISSUANCE OF THE EUA

The Secretary of Health and Human Services (HHS) has declared a public health emergency that justifies the emergency use of drugs and biological products during the COVID-19 pandemic. In response, FDA has issued an EUA for the unapproved product, Pfizer-BioNTech COVID-19 Vaccine, for active immunization against COVID-19 in individuals 12 years of age and older.

FDA issued this EUA, based on Pfizer-BioNTech's request and submitted data.

Although limited scientific information is available, based on the totality of the scientific evidence available to date, it is reasonable to believe that the Pfizer-BioNTech COVID-19 Vaccine may be effective for the prevention of COVID-19 in individuals as specified in the *Full EUA Prescribing Information*.

This EUA for the Pfizer-BioNTech COVID-19 Vaccine will end when the Secretary of HHS determines that the circumstances justifying the EUA no longer exist or when there is a change in the approval status of the product such that an EUA is no longer needed.

For additional information about Emergency Use Authorization visit FDA at: <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization>.

The Countermeasures Injury Compensation Program

The Countermeasures Injury Compensation Program (CICP) is a federal program that has been created to help pay for related costs of medical care and other specific expenses to compensate people injured after use of certain medical countermeasures. Medical countermeasures are specific vaccines, medications, devices, or other items used to prevent, diagnose, or treat the public during a public health emergency or a security threat. For more information about CICP regarding the Pfizer-BioNTech COVID-19 Vaccine used to prevent COVID-19, visit www.hrsa.gov/cicp, email cicp@hrsa.gov, or call: 1-855-266-2427.



Manufactured by
Pfizer Inc., New York, NY 10017

BIONTECH

Manufactured for
BioNTech Manufacturing GmbH
An der Goldgrube 12
55131 Mainz, Germany

LAB-1450-7.1

Revised: XX Month 2021

END SHORT VERSION FACT SHEET
Long Version (Full EUA Prescribing Information) Begins On Next Page

FULL EMERGENCY USE AUTHORIZATION (EUA) PRESCRIBING INFORMATION

PFIZER-BIONTECH COVID-19 VACCINE

FULL EMERGENCY USE AUTHORIZATION PRESCRIBING INFORMATION: CONTENTS*

- | | |
|--|---|
| <ul style="list-style-type: none">1 AUTHORIZED USE2 DOSAGE AND ADMINISTRATION<ul style="list-style-type: none">2.1 Preparation for Administration2.2 Administration Information2.3 Vaccination Schedule for Individuals 12 Years of Age and Older3 DOSAGE FORMS AND STRENGTHS4 CONTRAINDICATIONS5 WARNINGS AND PRECAUTIONS<ul style="list-style-type: none">5.1 Management of Acute Allergic Reactions5.2 Altered Immunocompetence5.3 Limitation of Effectiveness6 OVERALL SAFETY SUMMARY<ul style="list-style-type: none">6.1 Clinical Trials Experience6.2 Post Authorization Experience | <ul style="list-style-type: none">8 REQUIREMENTS AND INSTRUCTIONS FOR REPORTING
ADVERSE EVENTS AND VACCINE ADMINISTRATION
ERRORS10 DRUG INTERACTIONS11 USE IN SPECIFIC POPULATIONS<ul style="list-style-type: none">11.1 Pregnancy11.2 Lactation11.3 Pediatric Use11.4 Geriatric Use13 DESCRIPTION14 CLINICAL PHARMACOLOGY<ul style="list-style-type: none">14.1 Mechanism of Action18 CLINICAL TRIAL RESULTS AND SUPPORTING DATA FOR
EUA<ul style="list-style-type: none">18.1 Efficacy in Participants 16 Years of Age and Older18.2 Efficacy and Immunogenicity in Adolescents 12 to 15 Years of
Age19 HOW SUPPLIED/STORAGE AND HANDLING20 PATIENT COUNSELING INFORMATION21 CONTACT INFORMATION |
|--|---|

* Sections or subsections omitted from the full emergency use authorization prescribing information are not listed.

FULL EMERGENCY USE AUTHORIZATION (EUA) PRESCRIBING INFORMATION

1 AUTHORIZED USE

Pfizer-BioNTech COVID-19 Vaccine is authorized for use under an Emergency Use Authorization (EUA) for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 12 years of age and older.

2 DOSAGE AND ADMINISTRATION

For intramuscular injection only.

2.1 Preparation for Administration

Prior to Dilution

- The Pfizer-BioNTech COVID-19 Vaccine Multiple Dose Vial contains a volume of 0.45 mL, supplied as a frozen suspension that does not contain preservative. Each vial must be thawed and diluted prior to administration.
- Vials may be thawed in the refrigerator [2°C to 8°C (35°F to 46°F)] or at room temperature [up to 25°C (77°F)] [*see How Supplied/Storage and Handling (19)*].
- Refer to thawing instructions in the panels below.


Dilution

- Dilute the vial contents using 1.8 mL of 0.9% Sodium Chloride Injection, USP (not provided) to form the Pfizer-BioNTech COVID-19 Vaccine. Do not add more than 1.8 mL of diluent.
- ONLY use 0.9% Sodium Chloride Injection, USP as the diluent. This diluent is not packaged with the vaccine and must be sourced separately. Do not use bacteriostatic 0.9% Sodium Chloride Injection or any other diluent.
- After dilution, one vial contains 6 doses of 0.3 mL. Vial labels and cartons may state that after dilution, a vial contains 5 doses of 0.3 mL. The information in this Full EUA Prescribing Information regarding the number of doses per vial after dilution supersedes the number of doses stated on vial labels and cartons.
- Refer to dilution and dose preparation instructions in the panels below.

THAWING PRIOR TO DILUTION




- Thaw vial(s) of Pfizer-BioNTech COVID-19 Vaccine before use either by:
 - Allowing vial(s) to thaw in the refrigerator [2°C to 8°C (35°F to 46°F)]. A carton of vials may take up to 3 hours to thaw, and thawed vials can be stored in the refrigerator for up to five days (120 hours).
 - Allowing vial(s) to sit at room temperature [up to 25°C (77°F)] for 30 minutes.
- Using either thawing method, vials must reach room temperature before dilution and must be diluted within 2 hours.






- Before dilution invert vaccine vial gently 10 times.
- Do not shake.
- Inspect the liquid in the vial prior to dilution. The liquid is a white to off-white suspension and may contain white to off-white opaque amorphous particles.
- Do not use if liquid is discolored or if other particles are observed.


DILUTION



- Obtain sterile 0.9% Sodium Chloride Injection, USP. Use only this as the diluent.
- Using aseptic technique, withdraw 1.8 mL of diluent into a transfer syringe (21-gauge or narrower needle).
- Cleanse the vaccine vial stopper with a single-use antiseptic swab.
- Add 1.8 mL of 0.9% Sodium Chloride Injection, USP into the vaccine vial.

	<ul style="list-style-type: none"> Equalize vial pressure before removing the needle from the vial by withdrawing 1.8 mL air into the empty diluent syringe.
	<ul style="list-style-type: none"> Gently invert the vial containing the Pfizer-BioNTech COVID-19 Vaccine 10 times to mix. <u>Do not shake.</u> Inspect the vaccine in the vial. The vaccine will be an off-white suspension. Do not use if vaccine is discolored or contains particulate matter.
	<ul style="list-style-type: none"> Record the date and time of dilution on the Pfizer-BioNTech COVID-19 Vaccine vial label. Store between 2°C to 25°C (35°F to 77°F). Discard any unused vaccine 6 hours after dilution.

PREPARATION OF INDIVIDUAL 0.3 mL DOSES OF PFIZER-BIONTECH COVID-19 VACCINE



- Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab, and withdraw 0.3 mL of the Pfizer-BioNTech COVID-19 Vaccine preferentially using low dead-volume syringes and/or needles.
- Each dose must contain 0.3 mL of vaccine.
- If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and any excess volume.
- Administer immediately.

2.2 Administration Information

Visually inspect each dose in the dosing syringe prior to administration. The vaccine will be an off-white suspension. During the visual inspection,

- verify the final dosing volume of 0.3 mL.
- confirm there are no particulates and that no discoloration is observed.
- do not administer if vaccine is discolored or contains particulate matter.

Administer the Pfizer-BioNTech COVID-19 Vaccine intramuscularly.

After dilution, vials of Pfizer-BioNTech COVID-19 Vaccine contain six doses of 0.3 mL of vaccine. Low dead-volume syringes and/or needles can be used to extract six doses from a single vial. If standard syringes and needles are used, there may not be sufficient volume to extract a sixth dose from a single vial. Irrespective of the type of syringe and needle:

- Each dose must contain 0.3 mL of vaccine.
- If the amount of vaccine remaining in the vial cannot provide a full dose of 0.3 mL, discard the vial and any excess volume.
- Do not pool excess vaccine from multiple vials.

2.3 Vaccination Schedule for Individuals 12 Years of Age and Older

The Pfizer-BioNTech COVID-19 Vaccine is administered intramuscularly as a series of two doses (0.3 mL each) three weeks apart.

There are no data available on the interchangeability of the Pfizer-BioNTech COVID-19 Vaccine with other COVID-19 vaccines to complete the vaccination series. Individuals who have received one dose of Pfizer-BioNTech COVID-19 Vaccine should receive a second dose of Pfizer-BioNTech COVID-19 Vaccine to complete the vaccination series.

3 DOSAGE FORMS AND STRENGTHS

Pfizer-BioNTech COVID-19 Vaccine is a suspension for injection. After preparation, a single dose is 0.3 mL.

4 CONTRAINDICATIONS

Do not administer Pfizer-BioNTech COVID-19 Vaccine to individuals with known history of a severe allergic reaction (e.g., anaphylaxis) to any component of the Pfizer-BioNTech COVID-19 Vaccine [see *Description (13)*].

5 WARNINGS AND PRECAUTIONS

5.1 Management of Acute Allergic Reactions

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of Pfizer-BioNTech COVID-19 Vaccine.

Monitor Pfizer-BioNTech COVID-19 Vaccine recipients for the occurrence of immediate adverse reactions according to the Centers for Disease Control and Prevention guidelines (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-anaphylaxis.html>).

5.2 Altered Immunocompetence

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the Pfizer-BioNTech COVID-19 Vaccine.

5.3 Limitation of Effectiveness

The Pfizer-BioNTech COVID-19 Vaccine may not protect all vaccine recipients.

6 OVERALL SAFETY SUMMARY

It is MANDATORY for vaccination providers to report to the Vaccine Adverse Event Reporting System (VAERS) all vaccine administration errors, all serious adverse events, cases of Multisystem Inflammatory Syndrome (MIS) in adults and children, and hospitalized or fatal cases of COVID-19 following vaccination with the Pfizer-BioNTech COVID-19 Vaccine. To the extent feasible, provide a copy of the VAERS form to Pfizer Inc. Please see the REQUIREMENTS AND INSTRUCTIONS FOR REPORTING ADVERSE EVENTS AND VACCINE ADMINISTRATION ERRORS section for details on reporting to VAERS and Pfizer Inc.

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of Pfizer-BioNTech COVID-19 Vaccine was evaluated in participants 12 years of age and older in two clinical studies conducted in the United States, Europe, Turkey, South Africa, and South America. Study BNT162-01 (Study 1) was a Phase 1/2, two-part, dose-escalation trial that enrolled 60 participants, 18 through 55 years of age. Study C4591001 (Study 2) is a Phase 1/2/3, multicenter, multinational, randomized, saline placebo-controlled, observer-blind, dose-finding, vaccine candidate-selection (Phase 1) and efficacy (Phase 2/3) study that has enrolled approximately 46,000 participants, 12 years of age or older. Of these, approximately 43,448 participants (21,720 Pfizer-BioNTech COVID-19 Vaccine; 21,728 placebo) in Phase 2/3 are 16 years of age or older (including 138 and 145 adolescents 16 and 17 years of age in the vaccine and

placebo groups, respectively) and 2260 adolescents are 12 to 15 years of age (1131 and 1129 in the vaccine and placebo groups, respectively).

Participants 16 Years of Age and Older

At the time of the analysis of Study 2 for the EUA, 37,586 (18,801 Pfizer-BioNTech COVID-19 Vaccine and 18,785 placebo) participants 16 years of age or older had been followed for a median of 2 months after the second dose of Pfizer-BioNTech COVID-19 Vaccine.

The safety evaluation in Study 2 is ongoing. The safety population includes participants enrolled by October 9, 2020, and includes safety data accrued through November 14, 2020. Participants 18 years and older in the reactogenicity subset are monitored for solicited local and systemic reactions and use of antipyretic medication after each vaccination in an electronic diary. Participants are being monitored for unsolicited adverse events, including serious adverse events, throughout the study [from Dose 1 through 1 month (all unsolicited adverse events) or 6 months (serious adverse events) after the last vaccination].

Demographic characteristics in Study 2 were generally similar with regard to age, gender, race, and ethnicity among participants who received Pfizer-BioNTech COVID-19 Vaccine and those who received placebo. Overall, among the total participants who received either the Pfizer-BioNTech COVID-19 Vaccine or placebo, 50.6% were male and 49.4% were female, 83.1% were White, 9.1% were Black or African American, 28.0% were Hispanic/Latino, 4.3% were Asian, and 0.5% were American Indian/Alaska Native.

Local and Systemic Adverse Reactions Solicited in the Study 2

Table 1 and Table 2 present the frequency and severity of solicited local and systemic reactions, respectively, within 7 days following each dose of Pfizer-BioNTech COVID-19 Vaccine and placebo in the subset of participants 18 to 55 years of age included in the EUA safety population who were monitored for reactogenicity with an electronic diary.

Table 3 and Table 4 present the frequency and severity of reported solicited local and systemic reactions, respectively, within 7 days of each dose of Pfizer-BioNTech COVID-19 Vaccine and placebo for participants 56 years of age and older.

In clinical studies, adverse reactions in participants 16 years of age and older included pain at the injection site (84.1%), fatigue (62.9%), headache (55.1%), muscle pain (38.3%), chills (31.9%), joint pain (23.6%), fever (14.2%), injection site swelling (10.5%), injection site redness (9.5%), nausea (1.1%), malaise (0.5%), and lymphadenopathy (0.3%).

Across both age groups, the mean duration of pain at the injection site after Dose 2 was 2.5 days (range 1 to 36 days), for redness 2.6 days (range 1 to 34 days), and for swelling 2.3 days (range 1 to 34 days) for participants in the Pfizer-BioNTech COVID-19 Vaccine group.

Solicited reactogenicity data in 16 and 17 year-old participants are limited.

Table 1: Study 2 – Frequency and Percentages of Participants with Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 18-55 Years of Age[‡] – Reactogenicity Subset of the Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=2291 n^b (%)	Placebo Dose 1 N^a=2298 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=2098 n^b (%)	Placebo Dose 2 N^a=2103 n^b (%)
Redness^c				
Any (>2 cm)	104 (4.5)	26 (1.1)	123 (5.9)	14 (0.7)
Mild	70 (3.1)	16 (0.7)	73 (3.5)	8 (0.4)
Moderate	28 (1.2)	6 (0.3)	40 (1.9)	6 (0.3)
Severe	6 (0.3)	4 (0.2)	10 (0.5)	0 (0.0)
Swelling^c				
Any (>2 cm)	132 (5.8)	11 (0.5)	132 (6.3)	5 (0.2)
Mild	88 (3.8)	3 (0.1)	80 (3.8)	3 (0.1)
Moderate	39 (1.7)	5 (0.2)	45 (2.1)	2 (0.1)
Severe	5 (0.2)	3 (0.1)	7 (0.3)	0 (0.0)
Pain at the injection site^d				
Any	1904 (83.1)	322 (14.0)	1632 (77.8)	245 (11.7)
Mild	1170 (51.1)	308 (13.4)	1039 (49.5)	225 (10.7)
Moderate	710 (31.0)	12 (0.5)	568 (27.1)	20 (1.0)
Severe	24 (1.0)	2 (0.1)	25 (1.2)	0 (0.0)

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after vaccination.

a. N = Number of participants reporting at least 1 yes or no response for the specified reaction after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: >2.0 to ≤5.0 cm; Moderate: >5.0 to ≤10.0 cm; Severe: >10.0 cm.

d. Mild: does not interfere with activity; Moderate: interferes with activity; Severe: prevents daily activity.

[‡] Eight participants were between 16 and 17 years of age.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Table 2: Study 2 – Frequency and Percentages of Participants with Solicited Systemic Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 18-55 Years of Age[‡] – Reactogenicity Subset of the Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=2291 n^b (%)	Placebo Dose 1 N^a=2298 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=2098 n^b (%)	Placebo Dose 2 N^a=2103 n^b (%)
Fever				
≥38.0°C	85 (3.7)	20 (0.9)	331 (15.8)	10 (0.5)
≥38.0°C to 38.4°C	64 (2.8)	10 (0.4)	194 (9.2)	5 (0.2)
>38.4°C to 38.9°C	15 (0.7)	5 (0.2)	110 (5.2)	3 (0.1)
>38.9°C to 40.0°C	6 (0.3)	3 (0.1)	26 (1.2)	2 (0.1)
>40.0°C	0 (0.0)	2 (0.1)	1 (0.0)	0 (0.0)
Fatigue^c				
Any	1085 (47.4)	767 (33.4)	1247 (59.4)	479 (22.8)
Mild	597 (26.1)	467 (20.3)	442 (21.1)	248 (11.8)
Moderate	455 (19.9)	289 (12.6)	708 (33.7)	217 (10.3)
Severe	33 (1.4)	11 (0.5)	97 (4.6)	14 (0.7)

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=2291 n^b (%)	Placebo Dose 1 N^a=2298 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=2098 n^b (%)	Placebo Dose 2 N^a=2103 n^b (%)
Headache^c				
Any	959 (41.9)	775 (33.7)	1085 (51.7)	506 (24.1)
Mild	628 (27.4)	505 (22.0)	538 (25.6)	321 (15.3)
Moderate	308 (13.4)	251 (10.9)	480 (22.9)	170 (8.1)
Severe	23 (1.0)	19 (0.8)	67 (3.2)	15 (0.7)
Chills^c				
Any	321 (14.0)	146 (6.4)	737 (35.1)	79 (3.8)
Mild	230 (10.0)	111 (4.8)	359 (17.1)	65 (3.1)
Moderate	82 (3.6)	33 (1.4)	333 (15.9)	14 (0.7)
Severe	9 (0.4)	2 (0.1)	45 (2.1)	0 (0.0)
Vomiting^d				
Any	28 (1.2)	28 (1.2)	40 (1.9)	25 (1.2)
Mild	24 (1.0)	22 (1.0)	28 (1.3)	16 (0.8)
Moderate	4 (0.2)	5 (0.2)	8 (0.4)	9 (0.4)
Severe	0 (0.0)	1 (0.0)	4 (0.2)	0 (0.0)
Diarrhea^e				
Any	255 (11.1)	270 (11.7)	219 (10.4)	177 (8.4)
Mild	206 (9.0)	217 (9.4)	179 (8.5)	144 (6.8)
Moderate	46 (2.0)	52 (2.3)	36 (1.7)	32 (1.5)
Severe	3 (0.1)	1 (0.0)	4 (0.2)	1 (0.0)
New or worsened muscle pain^c				
Any	487 (21.3)	249 (10.8)	783 (37.3)	173 (8.2)
Mild	256 (11.2)	175 (7.6)	326 (15.5)	111 (5.3)
Moderate	218 (9.5)	72 (3.1)	410 (19.5)	59 (2.8)
Severe	13 (0.6)	2 (0.1)	47 (2.2)	3 (0.1)
New or worsened joint pain^c				
Any	251 (11.0)	138 (6.0)	459 (21.9)	109 (5.2)
Mild	147 (6.4)	95 (4.1)	205 (9.8)	54 (2.6)
Moderate	99 (4.3)	43 (1.9)	234 (11.2)	51 (2.4)
Severe	5 (0.2)	0 (0.0)	20 (1.0)	4 (0.2)
Use of antipyretic or pain medication^f	638 (27.8)	332 (14.4)	945 (45.0)	266 (12.6)

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose.

a. N = Number of participants reporting at least 1 yes or no response for the specified event after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: does not interfere with activity; Moderate: some interference with activity; Severe: prevents daily activity.

d. Mild: 1 to 2 times in 24 hours; Moderate: >2 times in 24 hours; Severe: requires intravenous hydration.

e. Mild: 2 to 3 loose stools in 24 hours; Moderate: 4 to 5 loose stools in 24 hours; Severe: 6 or more loose stools in 24 hours.

f. Severity was not collected for use of antipyretic or pain medication.

‡ Eight participants were between 16 and 17 years of age.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Table 3: Study 2 – Frequency and Percentages of Participants with Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 56 Years of Age and Older – Reactogenicity Subset of the Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=1802 n^b (%)	Placebo Dose 1 N^a=1792 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=1660 n^b (%)	Placebo Dose 2 N^a=1646 n^b (%)
Redness^c				
Any (>2 cm)	85 (4.7)	19 (1.1)	120 (7.2)	12 (0.7)
Mild	55 (3.1)	12 (0.7)	59 (3.6)	8 (0.5)
Moderate	27 (1.5)	5 (0.3)	53 (3.2)	3 (0.2)
Severe	3 (0.2)	2 (0.1)	8 (0.5)	1 (0.1)
Swelling^c				
Any (>2 cm)	118 (6.5)	21 (1.2)	124 (7.5)	11 (0.7)
Mild	71 (3.9)	10 (0.6)	68 (4.1)	5 (0.3)
Moderate	45 (2.5)	11 (0.6)	53 (3.2)	5 (0.3)
Severe	2 (0.1)	0 (0.0)	3 (0.2)	1 (0.1)
Pain at the injection site^d				
Any (>2 cm)	1282 (71.1)	166 (9.3)	1098 (66.1)	127 (7.7)
Mild	1008 (55.9)	160 (8.9)	792 (47.7)	125 (7.6)
Moderate	270 (15.0)	6 (0.3)	298 (18.0)	2 (0.1)
Severe	4 (0.2)	0 (0.0)	8 (0.5)	0 (0.0)

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after vaccination.

a. N = Number of participants reporting at least 1 yes or no response for the specified reaction after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: >2.0 to ≤5.0 cm; Moderate: >5.0 to ≤10.0 cm; Severe: >10.0 cm.

d. Mild: does not interfere with activity; Moderate: interferes with activity; Severe: prevents daily activity.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Table 4: Study 2 – Frequency and Percentages of Participants with Solicited Systemic Reactions, by Maximum Severity, Within 7 Days After Each Dose – Participants 56 Years of Age and Older – Reactogenicity Subset of the Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=1802 n^b (%)	Placebo Dose 1 N^a=1792 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=1660 n^b (%)	Placebo Dose 2 N^a=1646 n^b (%)
Fever				
≥38.0°C	26 (1.4)	7 (0.4)	181 (10.9)	4 (0.2)
≥38.0°C to 38.4°C	23 (1.3)	2 (0.1)	131 (7.9)	2 (0.1)
>38.4°C to 38.9°C	1 (0.1)	3 (0.2)	45 (2.7)	1 (0.1)
>38.9°C to 40.0°C	1 (0.1)	2 (0.1)	5 (0.3)	1 (0.1)
>40.0°C	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Fatigue^c				
Any	615 (34.1)	405 (22.6)	839 (50.5)	277 (16.8)
Mild	373 (20.7)	252 (14.1)	351 (21.1)	161 (9.8)
Moderate	240 (13.3)	150 (8.4)	442 (26.6)	114 (6.9)
Severe	2 (0.1)	3 (0.2)	46 (2.8)	2 (0.1)

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=1802 n^b (%)	Placebo Dose 1 N^a=1792 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=1660 n^b (%)	Placebo Dose 2 N^a=1646 n^b (%)
Headache^c				
Any	454 (25.2)	325 (18.1)	647 (39.0)	229 (13.9)
Mild	348 (19.3)	242 (13.5)	422 (25.4)	165 (10.0)
Moderate	104 (5.8)	80 (4.5)	216 (13.0)	60 (3.6)
Severe	2 (0.1)	3 (0.2)	9 (0.5)	4 (0.2)
Chills^c				
Any	113 (6.3)	57 (3.2)	377 (22.7)	46 (2.8)
Mild	87 (4.8)	40 (2.2)	199 (12.0)	35 (2.1)
Moderate	26 (1.4)	16 (0.9)	161 (9.7)	11 (0.7)
Severe	0 (0.0)	1 (0.1)	17 (1.0)	0 (0.0)
Vomiting^d				
Any	9 (0.5)	9 (0.5)	11 (0.7)	5 (0.3)
Mild	8 (0.4)	9 (0.5)	9 (0.5)	5 (0.3)
Moderate	1 (0.1)	0 (0.0)	1 (0.1)	0 (0.0)
Severe	0 (0.0)	0 (0.0)	1 (0.1)	0 (0.0)
Diarrhea^e				
Any	147 (8.2)	118 (6.6)	137 (8.3)	99 (6.0)
Mild	118 (6.5)	100 (5.6)	114 (6.9)	73 (4.4)
Moderate	26 (1.4)	17 (0.9)	21 (1.3)	22 (1.3)
Severe	3 (0.2)	1 (0.1)	2 (0.1)	4 (0.2)
New or worsened muscle pain^c				
Any	251 (13.9)	149 (8.3)	477 (28.7)	87 (5.3)
Mild	168 (9.3)	100 (5.6)	202 (12.2)	57 (3.5)
Moderate	82 (4.6)	46 (2.6)	259 (15.6)	29 (1.8)
Severe	1 (0.1)	3 (0.2)	16 (1.0)	1 (0.1)
New or worsened joint pain^c				
Any	155 (8.6)	109 (6.1)	313 (18.9)	61 (3.7)
Mild	101 (5.6)	68 (3.8)	161 (9.7)	35 (2.1)
Moderate	52 (2.9)	40 (2.2)	145 (8.7)	25 (1.5)
Severe	2 (0.1)	1 (0.1)	7 (0.4)	1 (0.1)
Use of antipyretic or pain medication	358 (19.9)	213 (11.9)	625 (37.7)	161 (9.8)

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose.

a. N = Number of participants reporting at least 1 yes or no response for the specified event after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: does not interfere with activity; Moderate: some interference with activity; Severe: prevents daily activity.

d. Mild: 1 to 2 times in 24 hours; Moderate: >2 times in 24 hours; Severe: requires intravenous hydration.

e. Mild: 2 to 3 loose stools in 24 hours; Moderate: 4 to 5 loose stools in 24 hours; Severe: 6 or more loose stools in 24 hours.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Unsolicited Adverse Events

Serious Adverse Events

In Study 2, among participants 16 to 55 years of age who had received at least 1 dose of vaccine or placebo (Pfizer-BioNTech COVID-19 Vaccine = 10,841; placebo = 10,851), serious adverse events from Dose 1 through up to 30 days after Dose 2 in ongoing follow-up were reported by 0.4% of Pfizer-BioNTech COVID-19 Vaccine recipients and by 0.3% of placebo recipients. In a similar analysis, in participants 56 years of age and older (Pfizer-BioNTech COVID-19 Vaccine = 7960, placebo = 7934), serious adverse events were reported by 0.8% of Pfizer-BioNTech COVID-19 Vaccine recipients and by 0.6% of placebo recipients who received at least 1 dose of Pfizer-BioNTech COVID-19 Vaccine or placebo, respectively. In these analyses, 91.6% of study participants had at least 30 days of follow-up after Dose 2.

Appendicitis was reported as a serious adverse event for 12 participants, and numerically higher in the vaccine group, 8 vaccine participants and 4 placebo participants. Currently available information is insufficient to determine a causal relationship with the vaccine. There were no other notable patterns or numerical imbalances between treatment groups for specific categories of serious adverse events (including neurologic, neuro-inflammatory, and thrombotic events) that would suggest a causal relationship to Pfizer-BioNTech COVID-19 Vaccine.

Non-Serious Adverse Events

In Study 2 in which 10,841 participants 16 to 55 years of age received Pfizer-BioNTech COVID-19 Vaccine and 10,851 participants received placebo, non-serious adverse events from Dose 1 through up to 30 days after Dose 2 in ongoing follow-up were reported in 29.3% of participants who received Pfizer-BioNTech COVID-19 Vaccine and 13.2% of participants in the placebo group, for participants who received at least 1 dose. Overall in a similar analysis in which 7960 participants 56 years of age and older received Pfizer-BioNTech COVID-19 Vaccine, non-serious adverse events within 30 days were reported in 23.8% of participants who received Pfizer-BioNTech COVID-19 Vaccine and 11.7% of participants in the placebo group, for participants who received at least 1 dose. In these analyses, 91.6% of study participants had at least 30 days of follow-up after Dose 2.

The higher frequency of reported unsolicited non-serious adverse events among Pfizer-BioNTech COVID-19 Vaccine recipients compared to placebo recipients was primarily attributed to local and systemic adverse events reported during the first 7 days following vaccination that are consistent with adverse reactions solicited among participants in the reactogenicity subset and presented in Tables 3 and 4. From Dose 1 through 30 days after Dose 2, reports of lymphadenopathy were imbalanced with notably more cases in the Pfizer-BioNTech COVID-19 Vaccine group (64) vs. the placebo group (6), which is plausibly related to vaccination. Throughout the safety follow-up period to date, Bell's palsy (facial paralysis) was reported by four participants in the Pfizer-BioNTech COVID-19 Vaccine group. Onset of facial paralysis was Day 37 after Dose 1 (participant did not receive Dose 2) and Days 3, 9, and 48 after Dose 2. No cases of Bell's palsy were reported in the placebo group. Currently available information is insufficient to determine a causal relationship with the vaccine. There were no other notable patterns or numerical imbalances between treatment groups for specific categories of non-serious adverse events (including other neurologic or neuro-inflammatory, and thrombotic events) that would suggest a causal relationship to Pfizer-BioNTech COVID-19 Vaccine.

Adolescents 12 to 15 Years of Age

In an analysis of Study 2, based on data up to the cutoff date of March 13, 2021, 2260 adolescents (1131 Pfizer-BioNTech COVID-19 Vaccine; 1129 placebo) were 12 to 15 years of age. Of these, 1308 (660 Pfizer-BioNTech COVID-19 Vaccine and 648 placebo) adolescents have been followed for at least

2 months after the second dose of Pfizer-BioNTech COVID-19 Vaccine. The safety evaluation in Study 2 is ongoing.

Adolescents are monitored for solicited local and systemic reactions and use of antipyretic medication after each vaccination in an electronic diary. Adolescents are being monitored for unsolicited adverse events, including serious adverse events, throughout the study [from Dose 1 through 1 month (all unsolicited adverse events) and 6 months (serious adverse events) after the last vaccination].

Demographic characteristics in Study 2 were generally similar with regard to age, gender, race, and ethnicity among adolescents who received Pfizer-BioNTech COVID-19 Vaccine and those who received placebo. Overall, among the adolescents who received the Pfizer-BioNTech COVID-19 Vaccine, 50.1% were male and 49.9% were female, 85.9% were White, 4.6% were Black or African American, 11.7% were Hispanic/Latino, 6.4% were Asian, and 0.4% were American Indian/Alaska Native.

Local and Systemic Adverse Reactions Solicited in the Study 2

Table 5 and Table 6 present the frequency and severity of solicited local and systemic reactions, respectively, within 7 days following each dose of Pfizer-BioNTech COVID-19 Vaccine and placebo in adolescents 12 to 15 years of age included in the EUA safety population who were monitored for reactogenicity with an electronic diary.

In clinical studies, adverse reactions in adolescents 12 to 15 years of age included pain at the injection site (90.5%), fatigue (77.5%), headache (75.5%), chills (49.2%), muscle pain (42.2%), fever (24.3%), joint pain (20.2%), injection site swelling (9.2%), injection site redness (8.6%), lymphadenopathy (0.8%), and nausea (0.4%).

The mean duration of pain at the injection site after Dose 1 was 2.4 days (range 1 to 10 days), for redness 2.4 days (range 1 to 16 days), and for swelling 1.9 days (range 1 to 5 days) for adolescents in the Pfizer-BioNTech COVID-19 Vaccine group.

Table 5: Study 2 – Frequency and Percentages of Adolescents With Solicited Local Reactions, by Maximum Severity, Within 7 Days After Each Dose – Adolescents 12 to 15 Years of Age – Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=1127 n^b (%)	Placebo Dose 1 N^a=1127 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=1097 n^b (%)	Placebo Dose 2 N^a=1078 n^b (%)
Redness^c				
Any (>2 cm)	65 (5.8)	12 (1.1)	55 (5.0)	10 (0.9)
Mild	44 (3.9)	11 (1.0)	29 (2.6)	8 (0.7)
Moderate	20 (1.8)	1 (0.1)	26 (2.4)	2 (0.2)
Severe	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Swelling^c				
Any (>2 cm)	78 (6.9)	11 (1.0)	54 (4.9)	6 (0.6)
Mild	55 (4.9)	9 (0.8)	36 (3.3)	4 (0.4)
Moderate	23 (2.0)	2 (0.2)	18 (1.6)	2 (0.2)
Severe	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=1127 n^b (%)	Placebo Dose 1 N^a=1127 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=1097 n^b (%)	Placebo Dose 2 N^a=1078 n^b (%)
Pain at the injection site^d				
Any	971 (86.2)	263 (23.3)	866 (78.9)	193 (17.9)
Mild	467 (41.4)	227 (20.1)	466 (42.5)	164 (15.2)
Moderate	493 (43.7)	36 (3.2)	393 (35.8)	29 (2.7)
Severe	11 (1.0)	0 (0.0)	7 (0.6)	0 (0.0)

Note: Reactions were collected in the electronic diary (e-diary) from Day 1 to Day 7 after vaccination.

a. N = Number of participants reporting at least 1 yes or no response for the specified reaction after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: >2.0 to ≤5.0 cm; Moderate: >5.0 to ≤10.0 cm; Severe: >10.0 cm.

d. Mild: does not interfere with activity; Moderate: interferes with activity; Severe: prevents daily activity.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Table 6: Study 2 – Frequency and Percentages of Adolescents with Solicited Systemic Reactions, by Maximum Severity, Within 7 Days After Each Dose – Adolescents 12 to 15 Years of Age – Safety Population*

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=1127 n^b (%)	Placebo Dose 1 N^a=1127 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=1097 n^b (%)	Placebo Dose 2 N^a=1078 n^b (%)
Fever				
≥38.0°C	114 (10.1)	12 (1.1)	215 (19.6)	7 (0.6)
≥38.0°C to 38.4°C	74 (6.6)	8 (0.7)	107 (9.8)	5 (0.5)
>38.4°C to 38.9°C	29 (2.6)	2 (0.2)	83 (7.6)	1 (0.1)
>38.9°C to 40.0°C	10 (0.9)	2 (0.2)	25 (2.3)	1 (0.1)
>40.0°C	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Fatigue^c				
Any	677 (60.1)	457 (40.6)	726 (66.2)	264 (24.5)
Mild	278 (24.7)	250 (22.2)	232 (21.1)	133 (12.3)
Moderate	384 (34.1)	199 (17.7)	468 (42.7)	127 (11.8)
Severe	15 (1.3)	8 (0.7)	26 (2.4)	4 (0.4)
Headache^c				
Any	623 (55.3)	396 (35.1)	708 (64.5)	263 (24.4)
Mild	361 (32.0)	256 (22.7)	302 (27.5)	169 (15.7)
Moderate	251 (22.3)	131 (11.6)	384 (35.0)	93 (8.6)
Severe	11 (1.0)	9 (0.8)	22 (2.0)	1 (0.1)
Chills^c				
Any	311 (27.6)	109 (9.7)	455 (41.5)	73 (6.8)
Mild	195 (17.3)	82 (7.3)	221 (20.1)	52 (4.8)
Moderate	111 (9.8)	25 (2.2)	214 (19.5)	21 (1.9)
Severe	5 (0.4)	2 (0.2)	20 (1.8)	0 (0.0)

	Pfizer-BioNTech COVID-19 Vaccine Dose 1 N^a=1127 n^b (%)	Placebo Dose 1 N^a=1127 n^b (%)	Pfizer-BioNTech COVID-19 Vaccine Dose 2 N^a=1097 n^b (%)	Placebo Dose 2 N^a=1078 n^b (%)
Vomiting^d				
Any	31 (2.8)	10 (0.9)	29 (2.6)	12 (1.1)
Mild	30 (2.7)	8 (0.7)	25 (2.3)	11 (1.0)
Moderate	0 (0.0)	2 (0.2)	4 (0.4)	1 (0.1)
Severe	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Diarrhea^c				
Any	90 (8.0)	82 (7.3)	65 (5.9)	43 (4.0)
Mild	77 (6.8)	72 (6.4)	59 (5.4)	38 (3.5)
Moderate	13 (1.2)	10 (0.9)	6 (0.5)	5 (0.5)
Severe	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
New or worsened muscle pain^c				
Any	272 (24.1)	148 (13.1)	355 (32.4)	90 (8.3)
Mild	125 (11.1)	88 (7.8)	152 (13.9)	51 (4.7)
Moderate	145 (12.9)	60 (5.3)	197 (18.0)	37 (3.4)
Severe	2 (0.2)	0 (0.0)	6 (0.5)	2 (0.2)
New or worsened joint pain^c				
Any	109 (9.7)	77 (6.8)	173 (15.8)	51 (4.7)
Mild	66 (5.9)	50 (4.4)	91 (8.3)	30 (2.8)
Moderate	42 (3.7)	27 (2.4)	78 (7.1)	21 (1.9)
Severe	1 (0.1)	0 (0.0)	4 (0.4)	0 (0.0)
Use of antipyretic or pain medication^f	413 (36.6)	111 (9.8)	557 (50.8)	95 (8.8)

Note: Events and use of antipyretic or pain medication were collected in the electronic diary (e-diary) from Day 1 to Day 7 after each dose.

a. N = Number of participants reporting at least 1 yes or no response for the specified event after the specified dose.

b. n = Number of participants with the specified reaction.

c. Mild: does not interfere with activity; Moderate: some interference with activity; Severe: prevents daily activity.

d. Mild: 1 to 2 times in 24 hours; Moderate: >2 times in 24 hours; Severe: requires intravenous hydration.

e. Mild: 2 to 3 loose stools in 24 hours; Moderate: 4 to 5 loose stools in 24 hours; Severe: 6 or more loose stools in 24 hours.

f. Severity was not collected for use of antipyretic or pain medication.

* Randomized participants in the safety analysis population who received at least 1 dose of the study intervention.

Unsolicited Adverse Events

In the analysis of Study 2 among adolescents 12 to 15 years of age, 98.3% of study participants had at least 30 days of follow-up after Dose 2 (1131 adolescents received Pfizer-BioNTech COVID-19 Vaccine and 1129 adolescents received placebo).

Serious Adverse Events

Serious adverse events were reported by 0.4% of Pfizer-BioNTech COVID-19 Vaccine recipients and by 0.2% of placebo recipients.

Non-Serious Adverse Events

Non-serious adverse events were reported by 5.7% of Pfizer-BioNTech COVID-19 Vaccine recipients and by 5.8% of placebo recipients.

6.2 Post Authorization Experience

The following adverse reactions have been identified during post authorization use of Pfizer-BioNTech COVID-19 Vaccine. Because these reactions are reported voluntarily, it is not always possible to reliably estimate their frequency or establish a causal relationship to vaccine exposure.

Immune System Disorders: severe allergic reactions, including anaphylaxis, and other hypersensitivity reactions (e.g., rash, pruritus, urticaria, angioedema)

Gastrointestinal Disorders: diarrhea, vomiting

Musculoskeletal and Connective Tissue Disorders: pain in extremity (arm)

8 REQUIREMENTS AND INSTRUCTIONS FOR REPORTING ADVERSE EVENTS AND VACCINE ADMINISTRATION ERRORS

See Overall Safety Summary (Section 6) for additional information.

The vaccination provider enrolled in the federal COVID-19 Vaccination Program is responsible for MANDATORY reporting of the listed events following Pfizer-BioNTech COVID-19 Vaccine to the Vaccine Adverse Event Reporting System (VAERS):

- Vaccine administration errors whether or not associated with an adverse event
- Serious adverse events* (irrespective of attribution to vaccination)
- Cases of Multisystem Inflammatory Syndrome (MIS) in children and adults
- Cases of COVID-19 that result in hospitalization or death

*Serious adverse events are defined as:

- Death
- A life-threatening adverse event
- Inpatient hospitalization or prolongation of existing hospitalization
- A persistent or significant incapacity or substantial disruption of the ability to conduct normal life functions
- A congenital anomaly/birth defect
- An important medical event that based on appropriate medical judgement may jeopardize the individual and may require medical or surgical intervention to prevent one of the outcomes listed above

Instructions for Reporting to VAERS

The vaccination provider enrolled in the federal COVID-19 Vaccination Program should complete and submit a VAERS form to FDA using one of the following methods:

- Complete and submit the report online: <https://vaers.hhs.gov/reportevent.html>, or
- If you are unable to submit this form electronically, you may fax it to VAERS at 1-877-721-0366. If you need additional help submitting a report you may call the VAERS toll-free information line at 1-800-822-7967 or send an email to info@vaers.org.

IMPORTANT: When reporting adverse events or vaccine administration errors to VAERS, please complete the entire form with detailed information. It is important that the information reported to FDA be as detailed and complete as possible. Information to include:

- Patient demographics (e.g., patient name, date of birth)
- Pertinent medical history
- Pertinent details regarding admission and course of illness
- Concomitant medications
- Timing of adverse event(s) in relationship to administration of the Pfizer-BioNTech COVID-19 Vaccine
- Pertinent laboratory and virology information
- Outcome of the event and any additional follow-up information if it is available at the time of the VAERS report. Subsequent reporting of follow-up information should be completed if additional details become available.

The following steps are highlighted to provide the necessary information for safety tracking:

1. In Box 17, provide information on Pfizer-BioNTech COVID-19 Vaccine and any other vaccines administered on the same day; and in Box 22, provide information on any other vaccines received within one month prior.
2. In Box 18, description of the event:
 - a. Write “Pfizer-BioNTech COVID-19 Vaccine EUA” as the first line.
 - b. Provide a detailed report of vaccine administration error and/or adverse event. It is important to provide detailed information regarding the patient and adverse event/medication error for ongoing safety evaluation of this unapproved vaccine. Please see information to include listed above.
3. Contact information:
 - a. In Box 13, provide the name and contact information of the prescribing healthcare provider or institutional designee who is responsible for the report.
 - b. In Box 14, provide the name and contact information of the best doctor/healthcare professional to contact about the adverse event.
 - c. In Box 15, provide the address of the facility where vaccine was given (NOT the healthcare provider’s office address).

Other Reporting Instructions

Vaccination providers may report to VAERS other adverse events that are not required to be reported using the contact information above.

To the extent feasible, report adverse events to Pfizer Inc. using the contact information below or by providing a copy of the VAERS form to Pfizer Inc.

Website	Fax number	Telephone number
www.pfizersafetyreporting.com	1-866-635-8337	1-800-438-1985

10 DRUG INTERACTIONS

There are no data to assess the concomitant administration of the Pfizer-BioNTech COVID-19 Vaccine with other vaccines.

11 USE IN SPECIFIC POPULATIONS

11.1 Pregnancy

Risk Summary

All pregnancies have a risk of birth defect, loss, or other adverse outcomes. In the US general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. Available data on Pfizer-BioNTech COVID-19 Vaccine administered to pregnant women are insufficient to inform vaccine-associated risks in pregnancy.

In a reproductive and developmental toxicity study, 0.06 mL of a vaccine formulation containing the same quantity of nucleoside-modified messenger ribonucleic acid (mRNA) (30 mcg) and other ingredients included in a single human dose of Pfizer-BioNTech COVID-19 Vaccine was administered to female rats by the intramuscular route on four occasions: 21 and 14 days prior to mating, and on gestation days 9 and 20. No vaccine-related adverse effects on female fertility, fetal development, or postnatal development were reported in the study.

11.2 Lactation

Risk Summary

Data are not available to assess the effects of Pfizer-BioNTech COVID-19 Vaccine on the breastfed infant or on milk production/excretion.

11.3 Pediatric Use

Emergency Use Authorization of Pfizer-BioNTech COVID-19 in adolescents 12 to 15 years of age is based on demonstration of safety, efficacy, and effectiveness. The effectiveness of the vaccine was demonstrated by comparison of the immune response in adolescents 12 to 15 years of age to participants 16 to 25 years of age. Emergency Use Authorization of Pfizer-BioNTech COVID-19 Vaccine in adolescents 16 and 17 years of age is based on extrapolation of safety and effectiveness from adults 18 years of age and older.

Emergency Use Authorization of Pfizer-BioNTech COVID-19 Vaccine does not include use in individuals younger than 12 years of age.

11.4 Geriatric Use

Clinical studies of Pfizer-BioNTech COVID-19 Vaccine include participants 65 years of age and older and their data contributes to the overall assessment of safety and efficacy [*see Overall Safety Summary (6.1) and Clinical Trial Results and Supporting Data for EUA (18.1)*]. Of the total number of Pfizer-BioNTech COVID-19 Vaccine recipients in Study 2 (N=20,033), 21.4% (n=4,294) were 65 years of age and older and 4.3% (n=860) were 75 years of age and older.

13 DESCRIPTION

The Pfizer-BioNTech COVID-19 Vaccine is supplied as a frozen suspension in multiple dose vials; each vial must be diluted with 1.8 mL of sterile 0.9% Sodium Chloride Injection, USP prior to use to form the vaccine. Each dose of the Pfizer-BioNTech COVID-19 Vaccine contains 30 mcg of a nucleoside-modified messenger RNA (modRNA) encoding the viral spike (S) glycoprotein of SARS-CoV-2.

Each dose of the Pfizer-BioNTech COVID-19 Vaccine also includes the following ingredients: lipids (0.43 mg (4-hydroxybutyl)azanediyl)bis(hexane-6,1-diyl)bis(2-hexyldecanoate), 0.05 mg 2[(polyethylene glycol)-2000]-N,N-ditetradecylacetamide, 0.09 mg 1,2-distearoyl-sn-glycero-3-phosphocholine, and 0.2 mg cholesterol), 0.01 mg potassium chloride, 0.01 mg monobasic potassium phosphate, 0.36 mg sodium chloride, 0.07 mg dibasic sodium phosphate dihydrate, and 6 mg sucrose. The diluent (0.9% Sodium Chloride Injection, USP) contributes an additional 2.16 mg sodium chloride per dose.

The Pfizer-BioNTech COVID-19 Vaccine does not contain preservative. The vial stoppers are not made with natural rubber latex.

14 CLINICAL PHARMACOLOGY

14.1 Mechanism of Action

The modRNA in the Pfizer-BioNTech COVID-19 Vaccine is formulated in lipid particles, which enable delivery of the RNA into host cells to allow expression of the SARS-CoV-2 S antigen. The vaccine elicits an immune response to the S antigen, which protects against COVID-19.

18 CLINICAL TRIAL RESULTS AND SUPPORTING DATA FOR EUA

18.1 Efficacy in Participants 16 Years of Age and Older

Study 2 is a multicenter, multinational, Phase 1/2/3, randomized, placebo-controlled, observer-blind, dose-finding, vaccine candidate-selection, and efficacy study in participants 12 years of age and older. Randomization was stratified by age: 12 through 15 years of age, 16 through 55 years of age, or 56 years of age and older, with a minimum of 40% of participants in the ≥ 56 -year stratum. The study excluded participants who were immunocompromised and those who had previous clinical or microbiological diagnosis of COVID-19. Participants with preexisting stable disease, defined as disease not requiring significant change in therapy or hospitalization for worsening disease during the 6 weeks before enrollment, were included as were participants with known stable infection with human immunodeficiency virus (HIV), hepatitis C virus (HCV), or hepatitis B virus (HBV).

In the Phase 2/3 portion of Study 2, based on data accrued through November 14, 2020, approximately 44,000 participants 12 years of age and older were randomized equally and received 2 doses of Pfizer-BioNTech COVID-19 Vaccine or placebo separated by 21 days. Participants are planned to be followed for up to 24 months, for assessments of safety and efficacy against COVID-19.

The population for the analysis of the primary efficacy endpoint included, 36,621 participants 12 years of age and older (18,242 in the Pfizer-BioNTech COVID-19 Vaccine group and 18,379 in the placebo group) who did not have evidence of prior infection with SARS-CoV-2 through 7 days after the second dose. Table 7 presents the specific demographic characteristics in the studied population.

Table 7: Demographics (population for the primary efficacy endpoint)^a

	Pfizer-BioNTech COVID-19 Vaccine (N=18,242) n (%)	Placebo (N=18,379) n (%)
Sex		
Male	9318 (51.1)	9225 (50.2)
Female	8924 (48.9)	9154 (49.8)

	Pfizer-BioNTech COVID-19 Vaccine (N=18,242) n (%)	Placebo (N=18,379) n (%)
Age (years)		
Mean (SD)	50.6 (15.70)	50.4 (15.81)
Median	52.0	52.0
Min, max	(12, 89)	(12, 91)
Age group		
≥12 through 15 years	46 (0.3)	42 (0.2)
≥16 through 17 years	66 (0.4)	68 (0.4)
≥16 through 64 years	14,216 (77.9)	14,299 (77.8)
≥65 through 74 years	3176 (17.4)	3226 (17.6)
≥75 years	804 (4.4)	812 (4.4)
Race		
White	15,110 (82.8)	15,301 (83.3)
Black or African American	1617 (8.9)	1617 (8.8)
American Indian or Alaska Native	118 (0.6)	106 (0.6)
Asian	815 (4.5)	810 (4.4)
Native Hawaiian or other Pacific Islander	48 (0.3)	29 (0.2)
Other ^b	534 (2.9)	516 (2.8)
Ethnicity		
Hispanic or Latino	4886 (26.8)	4857 (26.4)
Not Hispanic or Latino	13,253 (72.7)	13,412 (73.0)
Not reported	103 (0.6)	110 (0.6)
Comorbidities ^c		
Yes	8432 (46.2)	8450 (46.0)
No	9810 (53.8)	9929 (54.0)

- a. All eligible randomized participants who receive all vaccination(s) as randomized within the predefined window, have no other important protocol deviations as determined by the clinician, and have no evidence of SARS-CoV-2 infection prior to 7 days after Dose 2.
- b. Includes multiracial and not reported.
- c. Number of participants who have 1 or more comorbidities that increase the risk of severe COVID-19 disease
 - Chronic lung disease (e.g., emphysema and chronic bronchitis, idiopathic pulmonary fibrosis, and cystic fibrosis) or moderate to severe asthma
 - Significant cardiac disease (e.g., heart failure, coronary artery disease, congenital heart disease, cardiomyopathies, and pulmonary hypertension)
 - Obesity (body mass index ≥ 30 kg/m²)
 - Diabetes (Type 1, Type 2 or gestational)
 - Liver disease
 - Human Immunodeficiency Virus (HIV) infection (not included in the efficacy evaluation)

The population in the primary efficacy analysis included all participants 12 years of age and older who had been enrolled from July 27, 2020, and followed for the development of COVID-19 through November 14, 2020. Participants 18 to 55 years of age and 56 years of age and older began enrollment from July 27, 2020, 16 to 17 years of age began enrollment from September 16, 2020 and 12 to 15 years of age began enrollment from October 15, 2020.

The vaccine efficacy information is presented in Table 8.

Table 8: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Age Subgroup – Participants Without Evidence of Infection and Participants With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

First COVID-19 occurrence from 7 days after Dose 2 in participants without evidence of prior SARS-CoV-2 infection*			
Subgroup	Pfizer-BioNTech COVID-19 Vaccine N^a=18,198 Cases n1^b Surveillance Time^c (n2^d)	Placebo N^a=18,325 Cases n1^b Surveillance Time^c (n2^d)	Vaccine Efficacy % (95% CI)
All subjects ^e	8 2.214 (17,411)	162 2.222 (17,511)	95.0 (90.3, 97.6) ^f
16 to 64 years	7 1.706 (13,549)	143 1.710 (13,618)	95.1 (89.6, 98.1) ^g
65 years and older	1 0.508 (3848)	19 0.511 (3880)	94.7 (66.7, 99.9) ^g
First COVID-19 occurrence from 7 days after Dose 2 in participants with or without evidence of prior SARS-CoV-2 infection			
Subgroup	Pfizer-BioNTech COVID-19 Vaccine N^a=19,965 Cases n1^b Surveillance Time^c (n2^d)	Placebo N^a=20,172 Cases n1^b Surveillance Time^c (n2^d)	Vaccine Efficacy % (95% CI)
All subjects ^e	9 2.332 (18,559)	169 2.345 (18,708)	94.6 (89.9, 97.3) ^f
16 to 64 years	8 1.802 (14,501)	150 1.814 (14,627)	94.6 (89.1, 97.7) ^g
65 years and older	1 0.530 (4044)	19 0.532 (4067)	94.7 (66.8, 99.9) ^g

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhea; vomiting).

* Participants who had no evidence of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

- a. N = number of participants in the specified group.
- b. n1 = Number of participants meeting the endpoint definition.
- c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.
- d. n2 = Number of participants at risk for the endpoint.
- e. No confirmed cases were identified in adolescents 12 to 15 years of age.
- f. Credible interval for VE was calculated using a beta-binomial model with a beta (0.700102, 1) prior for $\theta = r(1-VE)/(1+r(1-VE))$, where r is the ratio of surveillance time in the active vaccine group over that in the placebo group.
- g. Confidence interval (CI) for vaccine efficacy is derived based on the Clopper and Pearson method adjusted to the surveillance time.

18.2 Efficacy and Immunogenicity in Adolescents 12 to 15 Years of Age

An analysis of Study 2 has been performed in adolescents 12 to 15 years of age up to a data cutoff date of March 13, 2021.

Efficacy

The vaccine efficacy information in adolescents 12 to 15 years of age presented in Table 9.

Table 9: Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2: Without Evidence of Infection and With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Blinded Placebo-Controlled Follow-up Period, Adolescents 12 to 15 Years of Age Evaluable Efficacy (7 Days) Population

First COVID-19 occurrence from 7 days after Dose 2 in adolescents 12 to 15 years of age without evidence of prior SARS-CoV-2 infection*			
	Pfizer-BioNTech COVID-19 Vaccine N^a=1005 Cases n^{1b} Surveillance Time^c (n^{2d})	Placebo N^a=978 Cases n^{1b} Surveillance Time^c (n^{2d})	Vaccine Efficacy % (95% CI^e)
Adolescents 12 to 15 Years of Age	0 0.154 (1001)	16 0.147 (972)	100.0 (75.3, 100.0)
First COVID-19 occurrence from 7 days after Dose 2 in adolescents 12 to 15 years of age with or without evidence of prior SARS-CoV-2 infection			
	Pfizer-BioNTech COVID-19 Vaccine N^a=1119 Cases n^{1b} Surveillance Time^c (n^{2d})	Placebo N^a=1110 Cases n^{1b} Surveillance Time^c (n^{2d})	Vaccine Efficacy % (95% CI^e)
Adolescents 12 to 15 Years of Age	0 0.170 (1109)	18 0.163 (1094)	100.0 (78.1, 100.0)

Note: Confirmed cases were determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR) and at least 1 symptom consistent with COVID-19 (symptoms included: fever; new or increased cough; new or increased shortness of breath; chills; new or increased muscle pain; new loss of taste or smell; sore throat; diarrhea; vomiting).

* Participants who had no evidence of past SARS-CoV-2 infection (i.e., N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

- a. N = Number of participants in the specified group.
- b. n1 = Number of participants meeting the endpoint definition.
- c. Total surveillance time in 1000 person-years for the given endpoint across all participants within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.
- d. n2 = Number of participants at risk for the endpoint.
- e. Confidence interval (CI) for vaccine efficacy is derived based on the Clopper and Pearson method adjusted for surveillance time.

Immunogenicity

In Study 2 an analysis of SARS-CoV-2 neutralizing titers in a randomly selected subset of participants was performed to demonstrate non-inferior immune responses (within 1.5-fold) comparing adolescents 12 to 15 years of age to participants 16 to 25 years of age who had no serological or virological evidence of past SARS-CoV-2 infection. The immune response to Pfizer-BioNTech COVID-19 Vaccine in adolescents 12 to 15 years of age (n=190) was noninferior to the immune response in participants 16 to 25 years of age (n=170), based on results for SARS-CoV-2 neutralizing titers at 1 month after Dose 2. The geometric mean titers (GMT) ratio of the adolescents 12 to 15 years of age group to the participants 16 to 25 years of age group was 1.76, with a 2-sided 95% CI of 1.47 to 2.10, meeting the 1.5-fold noninferiority criterion (the lower bound

of the 2-sided 95% CI for the geometric mean ratio [GMR] >0.67) which indicates a statistically greater response in the adolescents 12 to 15 years of age than that of participants 16 to 25 years of age.

19 HOW SUPPLIED/STORAGE AND HANDLING

Pfizer-BioNTech COVID-19 Vaccine Suspension for Intramuscular Injection, Multiple Dose Vials are supplied in a carton containing 25 multiple dose vials (NDC 59267-1000-3) or 195 multiple dose vials (NDC 59267-1000-2). After dilution, one vial contains 6 doses of 0.3 mL. Vial labels and cartons may state that after dilution, a vial contains 5 doses of 0.3 mL. The information in this Full EUA Prescribing Information regarding the number of doses per vial after dilution supersedes the number of doses stated on vial labels and cartons.

During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light.

Do not refreeze thawed vials.

Frozen Vials Prior to Use

Cartons of Pfizer-BioNTech COVID-19 Vaccine Multiple Dose Vials arrive in thermal containers with dry ice. Once received, remove the vial cartons immediately from the thermal container and preferably store in an ultra-low temperature freezer between -80°C to -60°C (-112°F to -76°F) until the expiry date printed on the label. Alternatively, vials may be stored at -25°C to -15°C (-13°F to 5°F) for up to 2 weeks. Vials must be kept frozen and protected from light, in the original cartons, until ready to use. Vials stored at -25°C to -15°C (-13°F to 5°F) for up to 2 weeks may be returned one time to the recommended storage condition of -80°C to -60°C (-112°F to -76°F). Total cumulative time the vials are stored at -25°C to -15°C (-13°F to 5°F) should be tracked and should not exceed 2 weeks.

If an ultra-low temperature freezer is not available, the thermal container in which the Pfizer-BioNTech COVID-19 Vaccine arrives may be used as temporary storage when consistently re-filled to the top of the container with dry ice. Refer to the re-icing guidelines packed in the original thermal container for instructions regarding the use of the thermal container for temporary storage. The thermal container maintains a temperature range of -90°C to -60°C (-130°F to -76°F). Storage of the vials between -96°C to -60°C (-141°F to -76°F) is not considered an excursion from the recommended storage condition.

Transportation of Frozen Vials

If local redistribution is needed and full cartons containing vials cannot be transported at -90°C to -60°C (-130°F to -76°F), vials may be transported at -25°C to -15°C (-13°F to 5°F). Any hours used for transport at -25°C to -15°C (-13°F to 5°F) count against the 2-week limit for storage at -25°C to -15°C (-13°F to 5°F). Frozen vials transported at -25°C to -15°C (-13°F to 5°F) may be returned one time to the recommended storage condition of -80°C to -60°C (-112°F to -76°F).

Thawed Vials Before Dilution

Thawed Under Refrigeration

Thaw and then store undiluted vials in the refrigerator [2°C to 8°C (35°F to 46°F)] for up to 5 days (120 hours). A carton of 25 vials or 195 vials may take up to 2 or 3 hours, respectively, to thaw in the refrigerator, whereas a fewer number of vials will thaw in less time.

Thawed at Room Temperature

For immediate use, thaw undiluted vials at room temperature [up to 25°C (77°F)] for 30 minutes. Thawed vials can be handled in room light conditions.

Vials must reach room temperature before dilution.

Undiluted vials may be stored at room temperature for no more than 2 hours.

Transportation of Thawed Vials

Available data support transportation of one or more thawed vials at 2°C to 8°C (35°F to 46°F) for up to 12 hours. Any hours used for transport at 2°C to 8°C (35°F to 46°F) count against the 120-hour limit for storage at 2°C to 8°C (35°F to 46°F).

Vials After Dilution

After dilution, store vials between 2°C to 25°C (35°F to 77°F) and use within 6 hours from the time of dilution. During storage, minimize exposure to room light, and avoid exposure to direct sunlight and ultraviolet light. Any vaccine remaining in vials must be discarded after 6 hours. Do not refreeze.


20 PATIENT COUNSELING INFORMATION

Advise the recipient or caregiver to read the Fact Sheet for Recipients and Caregivers.

The vaccination provider must include vaccination information in the state/local jurisdiction's Immunization Information System (IIS) or other designated system. Advise recipient or caregiver that more information about IISs can be found at: <https://www.cdc.gov/vaccines/programs/iis/about.html>.

21 CONTACT INFORMATION

For general questions, visit the website or call the telephone number provided below.

Website	Telephone number
www.cvdvaccine.com 	1-877-829-2619 (1-877-VAX-CO19)

This Full EUA Prescribing Information may have been updated. For the most recent Full EUA Prescribing Information, please see www.cvdvaccine.com.



Manufactured by
Pfizer Inc., New York, NY 10017

BIONTECH

Manufactured for
BioNTech Manufacturing GmbH
An der Goldgrube 12
55131 Mainz, Germany

LAB-1457-7.1

Revised: XX Month 2021